

ON THE STRUCTURE OF GENERAL SOLUTION TO THE EQUATIONS OF SHEAR-FREE NULL CONGRUENCES

Vladimir V. Kassandrov

(Institute of Gravitation and Cosmology,
Russian People's Friendship University, Moscow, Russia)

Abstract. We demonstrate that the source of a shear-free null congruence is, generically, a world sheet of a singular string in the complex Minkowski space. In particular cases the string degenerates into the Newman's point singularity. We describe also a time-dependent deformation of the Kerr congruence which could indicate the instability of the Kerr (Kerr-Newman) (electro)vacuum solution. Then we consider a 5D extension of Minkowski space obtained by complexification of the time coordinate. Singular locus of a shear-free null congruence on such an extended space-time consists of a (great) number of point particles ("markeons") and 2D surfaces – light-like wave fronts.

Congruences of light rays, with zero shear in particular, play crucial role in general relativity and in twistor aspects of geometry of flat Minkowski space-time $\mathbf{M}$. They relate also to the conditions of differentiability for the biquaternion-valued functions which may be put into correspondence with fundamental physical fields (in the framework of the so called *algebrodynamical* approach, see below).

General solution to the defining equations of shear-free (null geodesic) congruences (SFC) is represented by the *Kerr theorem* and can be formulated in the following implicit algebraic form [1]:

$$\Pi(\xi, iX\xi) = 0, \quad (1)$$

where $\Pi(\xi, \tau)$ is an *arbitrary homogeneous* (and almost everywhere analytical) function of three *projective* components of the twistor $\{\xi, \tau\}$ on $\mathbf{M}$, i.e. of a pair of the 2-spinors linked at every space-time point $X = X^+ = \{X_{AA'}\}$, $A = A' = 0, 1$ via the Penrose *incidence relation*,

$$\tau = iX\xi \quad (\tau^A = iX^{AA'}\xi_{A'}) \quad (2)$$

(see, e.g., [1], ch 7). At a fixed point X , resolving (1) with respect to the ratio of components of the spinor $\xi_A = \xi_A(X)$, one comes to a null 4-vector field $k_\mu = \xi_A \xi_{A'}$, $\mu = 0, 1, 2, 3$ tangent to the congruence which is necessarily shear-free; according to the Kerr theorem, all (analytical) SFC can be constructed via the above presented algebraic procedure.

Caustics - singularities of SFC - are defined by the condition

$$\Pi'(\xi) = 0, \quad (3)$$

i.e. correspond to the space-time points at which the *total derivative* of Π with respect to the *ratio* of spinor components turns to zero. Eliminating the latter from eq.(3) and substituting the result into eq.(1), one comes to the *equation of motion*

$$S(X) \equiv \Pi(\xi(X), iX\xi(X)) = 0 \quad (4)$$

which defines the shape of singular locus of the congruence (at a fixed moment of time) and its time evolution as well. It has been shown [3, 4] that any such function $S(X)$ satisfies the *complex eikonal equation* (CEE).

In the framework of *algebrodynamics* (see [2, 3, 4] and references therein) such singularities (when bounded in 3-space) were identified with *particle-like* formations. A remarkable example of these in general relativity is represented by the Kerr's *singular ring* – the caustic of the twofold static and axisymmetrical Kerr congruence. It possesses some of the quantum numbers of elementary fermion (Dirac electron), see, e.g. [5]. Newman et al. [6, 7] have shown that the Kerr-like congruences may be viewed as being generated by a “virtual” point charge moving along a complex world line $Z_\mu = Z_\mu(\sigma)$, $\sigma \in \mathbb{C}$ in the *complex extension* $\mathbb{CM}$ of Minkowski space-time $\mathbf{M}$ and therewith emanating rectilinear complex “light-like rays”. Restriction of such a complex congruence to the real $\mathbf{M}$ gives there rise to a SFC, to the Kerr one in particular (the latter in the case of virtual charge at rest).

However, this construction [8] covers only a narrow subclass of SFC. Below we present the theorem which specifies the source of a SFC of a *generic type*. Precisely, we define the source of a SFC as the *hyper-caustic* set, i.e. the singular locus at which, apart from the conditions (1) and (3), the *second* total derivative (with respect to the ratio of spinor components) turns to zero,

$$\Pi''(\xi) = 0, \quad (5)$$

so that one deals here with the singularities of caustics – the *cusps*. The theorem can be formulated as follows.

Theorem.

The source of a shear-free congruence of a generic type is a *complex string* $Z_\mu = Z_\mu(\lambda, \sigma)$, $\lambda, \sigma \in \mathbb{C}$, i. e. a 1-dimensional complex curve $Z_\mu = Z_\mu(\lambda)$ evolving in “complex time” σ . This is determined by the set of algebraic eqs. (1), (3), (5). Precisely, at any point $Z \in \mathbb{CM}$ of complex Minkowski space the value of (projective) twistor $\{\xi, \tau\}$ of a SFC is the same as at some point of the generating string $Z_\mu = Z_\mu(\lambda, \sigma)$, and correspondent point belongs to the complex light cone of the string element.

The proof is very simple. Indeed, system of the *three* eqs.(1), (3), (5) together with the *two* incidence constraints (2), in all of which in the considered case of full complex space $\mathbb{CM}$ the substitution of coordinates $iX \mapsto Z$ is made, determine, generically, *five* of the *seven* unknowns – *four* complex coordinates Z^{AB} and *three* projective twistor components. In result, as a solution of the joint system one obtains

$$Z_\mu = Z_\mu(\tau^1, \tau^2), \quad (6)$$

where some two twistor components, say τ^A , are taken to parametrize the world sheet of the complex string whereas the third one, say $G = \xi_2/\xi_1$, is then defined by the basic Kerr constraint (1). Of course, generally any other parametrization of the world sheet could be chosen.

Let now a “branch” of the (generally multi-valued) projective twistor field $\{\xi, \tau\}$ of a SFC is obtained at a point $Z \in \mathbb{CM}$ as a solution of the basic Kerr eq. (1). Then eq. (6)

immediately specifies an element of generating string at which all of the components of the twistor field take the same values as at the considered point Z . $\square$

In other words, if a *world-sheet* $Z_\mu(\lambda, \sigma)$ of complex string is known, the congruence is easily reconstructed both on $\mathbb{CM}$ and $\mathbf{M}$ via examination of rectilinear complex null “rays” emanating by all its elements, and any SFC can be obtained in this way. In particular cases (e.g., for axisymmetrical congruences) the source string degenerates into the Newman’s point charge. Moreover, for some types of SFC the world sheet of generating string can *coexist* with the world line of the Newman’s point charge – *focal line* of the congruence. Examples of such congruences, perhaps the most interesting ones, will be presented elsewhere.

Simple considerations on codimensions show that, generically, the *caustic* locus on $\mathbf{M}$ consists of a number of *real isolated strings*, usually non-bounded and unstable in time. Axisymmetrical Kerr’s SFC with the Kerr’s singular ring is only an exceptional example of a bounded and “nonradiating” caustic. Indeed, by continuous change of parameters of the Kerr’s generating function Π it can be deformed to the form

$$\Pi = \tau^2(1 + ih) - G\tau^1(1 - ih) + 2iaG, \quad (7)$$

with a new parameter $h \in \mathbb{R}$ and the parameter $a \in \mathbb{R}$ being the old Kerr’s one. Then one obtains a time-dependent axisymmetrical congruence generated by the Newman’s charge moving uniformly in $\mathbb{CM}$ with “imaginary velocity” $v = ih$. Caustic locus of correspondent SFC in $\mathbf{M}$ is now represented by an uniformly collapsing and then expanding ring

$$z = 0, \quad \rho = \sqrt{x^2 + y^2} = v(t - t_0), \quad (8)$$

where $t_0 = a/\sqrt{1 + h^2}$ and the speed $v = h/\sqrt{1 + h^2}$ of contraction (at $t < t_0$) or expansion (at $t > t_0$) is always less than the speed of light $c = 1$. In account of existence of such a deformation of the Kerr congruence one can even suspect that the very *Kerr (Kerr-Newman) solution to Einstein (Einstein-Maxwell) (electro)vacuum system is unstable* as its generating SFC does.

Consider now the *hyper-caustic* locus of a generic SFC on $\mathbf{M}$. It is easy to see, in account of similar considerations on codimension, that it will be represented by a system of isolated points – *instantons*¹ – which appear only at some particular instants and disappear immediately.

All this seems to be far from the observable structure of physical matter. There is, however, a successive hypothesis which drastically changes the situation. Namely, let us consider a *five-dimensional* submanifold of $\mathbb{CM}$ formed via restriction onto $\mathbf{M}$ of three space-like coordinates, $Z_a \rightarrow X_a \in \mathbb{R}$, $a = 1, 2, 3$, whereas the time-like coordinate $Z_0 = t + is$, $t, s \in \mathbb{R}$ is assumed to remain essentially complex-valued. On such a 5D cut of $\mathbb{CM}$ singular locus is represented by the 2D surfaces which evolve in *real* time t , whereas the structure of hyper-caustics – by a (great) number of point charges – *markeons* (from English “mark”) – moving along their world lines, effectively interacting each other and stable by definition. As to the singular 2D surfaces, they strongly resemble the *light-like wave fronts*.

¹This term here has, certainly, nothing to do with the familiar Yang-Mills instantons

Thus, we see that assumption of the 5D structure of physical space-time (formed by complexification of the time-like parameter) leads to a trustful structure of elementary matter with the point-like stable “markeons” and the singular wave fronts as the basic constituents. Moreover, it can be easily demonstrated that the complex null cone (CNC) on such a 5D subspace *degenerates into the familiar light cone of the 4D Minkowski space-time*. Indeed, the CNC equation

$$(Z_0)^2 - (Z_1)^2 - (Z_2)^2 - (Z_3)^2 = 0 \quad (9)$$

for the case of three real and one complex time-like coordinate corresponds to two *real* constraints from the second of which $2ts = 0$ one gets $s = 0$ so that the first constraint reduces to the real Minkowski light cone

$$t^2 - x^2 - y^2 - z^2 = 0. \quad (10)$$

On the other hand, this means that the fifth coordinate s *remains constant along any ray of the congruence* and does not contribute, therefore, into the observable metrics of physical space-time. In other words, reduction of the extended space to the 4D physical one takes here place *dynamically*, i.e. despite any known artificial assumptions like cylindricity condition, compactification etc. Dynamics of markeons and their interaction with singular wave fronts are under study.

References

- [1] R. Penrose, W. Rindler, *Spinors and Space-Time. Vol.II. Spinor and Twistor Methods in Space-Time Geometry*. Cambridge Univ. Press, Cambridge, 1986.
- [2] V.V. Kassandrov, *Algebraic Structure of Space-Time and Algebrodynamics*. People’s Friend. Univ. Press, Moscow, 1992 (in Russian).
- [3] V.V. Kassandrov, // *Hypercomplex Numbers in Geom. & Phys.*, **1(1)**, 91-107, (2004); (hep-th / 0312278).
- [4] V.V. Kassandrov, // *Grav. & Cosm.*, **8**, Suppl. 2, 57-62, (2002); (math-ph / 0311006).
- [5] A.Ya. Burinskii, // *Phys. Rev.*, **D67**, 12024, (2003); (gr-qc/0212048).
- [6] R.W. Lind, E.T. Newman, // *Journ. Math. Phys.*, **15**, 1103-1114, (1974).
- [7] E.T. Newman, // *Phys. Rev.*, **D65**, 104005, (2002); (gr-qc / 0201055).
- [8] E.T. Newman // *Class. Quant. Grav.*, **21**, 3197-3222 (2004); (gr-qc / 0402056).